

- mill vs over-ground training for walking after acute incomplete SCI[J].Neurology,2006,66(4):484—493.
- [5] Leahy TE. Impact of a limited trial of walking training using body weight support and a treadmill on the gait characteristics of an individual with chronic, incomplete spinal cord injury[J].Physiother Theory Pract,2010,26(7):483—489.
- [6] 郭素梅,李建民,吴庆文,等. Lokomat全自动机器人步态训练与评定系统对不完全性脊髓损伤患者步行功能的影响[J]. 中国组织工程研究,2012,16,(13):2324—2325.
- [7] Mehrholz J,Kugler J,Pohl M.Locomotor training for walking after spinal cord injury[J].Spine(Phlla Pa 1976),2008,33E68—777.
- [8] Colombo G, Wirz M, Dietz V. Driven gait orthosis for improvement of locomotor training in paraplegia patients[J]. Spinal Cord, 2001, 39, 252—255.
- [9] Field-Fote EC, Roach KE. Influence of a locomotor training approach on walking speed and distance in people with chronic spinal cord injury: a randomized clinical trial[J]. Physical Therapy, 2011,91(1): 1—13.
- [10] Wellner M, Thüring T, Smajic E, et al. Obstacle crossing in a virtual environment with the rehabilitation gait robot Lokomat[J]. Stud Health Technol Inform,2007,125:497—499.
- [11] Lünenburger L, Colombo G, Riener R, et al. Biofeedback in gait training with the robotic orthosis Lokomat[J]. Conf Proc IEEE Eng Med Biol Soc, 2004,7:4888—4891.
- [12] Nash MS, Jacobs PL, Johnson BM, et al. metabolic and cardiac responses to robotic-assisted locomotion in motor-complete tetraplegia: a case report[J]. J Spinal Cord Med, 2004,27(1):78—82.
- [13] Hornby TG, Zemon DH, Campbell D. Robotic-assisted, body-weight-supported treadmill training in individuals following motor incomplete spinal cord injury[J]. Phys Ther, 2005,85(1):52—66.
- [14] Wirz M, Zemon DH, Rupp R, et al. Effectiveness of automated locomotor training in patients with chronic incomplete spinal cord injury: a multicenter trial[J]. Arch Phys Med Rehabil, 2005,86(4):672—680.
- [15] Nooijen CF, Ter Hoeve N, Field-Fote EC. Gait quality is improved by locomotor training in individuals with SCI regardless of training approach[J]. J Neuroeng Rehabil,2009,6: 36.
- [16] Field-Fote EC, Lindley SD, Sherman AL. Locomotor training approaches for individuals with spinal cord injury: a preliminary report of walking-related outcomes[J]. J Neurol Phys Ther,2005,29(3):127—137.
- [17] Mayr A, Kofler M, Quirbach E, et al. Prospective, blinded, randomized crossover study of gait rehabilitation in stroke patients using the Lokomat gait orthosis[J]. Neurorehabil Neural Repair,2007,21(4):307—314.
- [18] Westlake KP, Patten C. Pilot study of Lokomat versus manual-assisted. treadmill training for locomotor recovery post-stroke[J]. J Neuroeng Rehabil, 2009,6:18.
- [19] Hidler J, Nichols D, Pelliccio M, et al. Multicenter randomized clinical trial evaluating the effectiveness of the Lokomat in subacute stroke[J]. Neurorehabil Neural Repair, 2009, 23(1):5—13.
- [20] Hornby TG, Campbell DD, Kahn JH, et al. Enhanced gait-related improvements after therapist- versus robotic-assisted locomotor training in subjects with chronic stroke: a randomized controlled study[J]. Stroke,2008,39(6):1786—1792.

2018年北京大学第一医院举办国家级继教学习班通知

Alberta 婴儿运动量表及高危儿早期干预培训班 暨 Peabody 发育评估与干预方案培训班

内容:①Alberta 婴儿运动量表是一个通过观察来评估0—18个月龄婴儿运动发育的工具,与以往经典的里程碑式的运动发育量表相比,它更注重对婴儿的运动质量的评估,因此可以较早地识别运动发育不成熟或运动模式异常的婴儿,适用于高危儿早期监测,并为干预方案的制定提供有价值的参考信息;②以2016年翻译出版的《婴幼儿期脑性瘫痪:目标性活动优化早期生长和发育》为教材,讲授高危儿或脑瘫早期干预思路和技术要点。该书以大量研究成果为依据,结合儿童发育特点,从不同角度,不同层面,全面阐述了脑瘫的特征及其形成原因,提出了以终为始的早期干预策略和技术,剖析了有针对性设计的目标性活动训练对优化高危儿发育的重要性。培训班主讲教师由黄真主任医师及部分参译专家承担,将紧密联系临床,强调实用性。授课内容不仅有助于治疗师学习规范的评定和治疗技术,更有助于提高康复医师临床思维能力和临床技能。时间:Alberta 培训班2018年4月26—28日(25日报到),紧接在往年Peabody 培训班之后;Peabody 培训班2018年4月22—25日(21日报到)。参加两个培训班者获两个国家级继续教育学分。食宿统一安排,费用自理。报名方式:Alberta 学习班报名可电子邮件联系王翠:wanguibill@sina.com,也可电话联系:13811093176;Peabody 学习班报名可电子邮件联系田甜:cnhc_pku@aliyun.com,也可电话联系:18710030337。

北京大学第一医院